中国外来入侵植物的学名考证(3)*

寿海洋, 闫小玲, 马金双**

(中国科学院上海辰山植物科学研究中心/上海辰山植物园,上海 201602)

摘要:基于中国外来入侵植物的相关报道与文献,发现一些入侵植物的学名使用比较混乱:不同时期、不同地区、不同作者在报道同一外来物种时使用不同的学名,或因中文名相近,而将其他植物作为外来入侵植物报道。依据最新资料,对国内报道的 16 种外来入侵植物学名进行了考证,包括中文名、英文名、异名、原产地及其在国内外的分布情况。

关键词:中国:外来:入侵植物:学名考证

中图分类号: 0 949, 0 16

文献标识码: A

文章编号: 2095-0845(2014)06-763-08

Nomenclatural Notes on Alien Invasive Vascular Plants in China (3)*

SHOU Hai-Yang, YAN Xiao-Ling, MA Jin-Shuang**

(Shanghai Chenshan Plant Science Research Center, Chines Academy of Sciences, Shanghai Chenshan Botanical Garden, Shanghai 201602, China)

Abstract: Based on literatures of alien invasive plants in China, the scientific names of sixteen species are revised, since their records in the Chinese literatures are confused: the same species with different scientic names were used by different authors in different regions and in different times, even different species with similar Chinese names were treated as identical. The data from floras updated were emplyed to compile the scientific name of each species, along with vernacular name, synonym, their origin and distribution both in China and abroad.

Key words: China; Alien; Invasive plants; Nomenclatural notes

Amaranthus cruentus Linnaeus, Syst. Nat. (ed. 10)
 12: 1269. 1759; Bao B. J., S. E. Clemants & T. Borsch in Fl. Chin.
 1419. 2003; S. L. Mosyakin in Fl. N. Amer.
 14: 410. 2003. Syn. Amaranthus paniculatus Linnaeus, Sp. Pl. (ed. 2)
 1406. 1763 (Amaranthaceae).

Blood amaranth (老鸦谷), cultivated as ornamental and pseudocereal almost worldwide from tropical to warm-temperate regions, and sometimes escaped and naturalized near places of cultivation, has

been widely cultivated in China: Anhui, Beijing, Chongqing, Fujian, Gansu, Guangdong, Guangxi, Guizhou, Hebei, Henan, Heilongjiang, Hubei, Hunan, Jiangsu, Jilin, Jiangxi, Liaoning, Nei Mongol, Qinghai, Shaanxi, Shandong, Shanxi, Shanghai, Sichuan, Tianjin, Xizang, Yunnan, Zhejiang^①. It usually called as fan sui xian (繁穗克) in Chinese but treated as *Amaranthus paniculatus* (曲波, 2003; Qu et al., 2006a,b; Li et al., 2007; Wang et al., 2007; Fu et al., 2009; Gao and Cao, 2010; Chen

^{*} Funding: Shanghai Landscaping Administrative Bureau (G1024011)

This is our last report on the series, and two previous reports could be found in *Plant Diversity and Resources* (植物分类与资源学报) **33** (1): 132-142, 2011 and **34** (4): 347-353, 2012

^{**} Author for correspondence; E-mail: jinshuangma@gmail.com Received date: 2014-03-14, Accepted date: 2014-06-03

作者简介:寿海洋 (1983-) 男,硕士,工程师,主要从事种子植物分类学及植物资源学方面的研究。E-mail: shouhy@126.com

① According to The Checklist of the Chinese Invasive Plants (Ma, 2013, same as below)

et al., 2011; 万方浩等, 2012; Fu et al., 2012; Lin et al., 2012a; Wang et al., 2012).

Bromus catharticus Vahl, Symb. Bot. 2: 22. 1791.
 Liu L., Zhu G. H. & K. H. Ammann in Fl. Chin.
 386. 2006; Hsu C.-C. in Fl. Taiwan (2nd Edition)
 370. 2000. Syn. Bromus unioloides Kunth,
 Nov. Gen. Sp. 1: 151. 1815 (Gramineae).

Rescuegrass (扁穗雀麦), native to South America, widely introduced as a winter forage now has been adventive in most temperate countries of Asia, Europe, Australia, North America and China; Beijing, Gansu, Guangdong, Guangxi, Guizhou, Hebei, Hubei, Jiangsu, Nei Mongol, Qinghai, Shaanxi, Sichuan, Taiwan, Xinjiang, Yunnan. In some literatures, it's treated as *Bromus unioloides* (Wu et al., 2004).

3. Crotalaria micans Link, Enum. Hort. Berol. Alt. 2: 228-229. 1822. Li J. Q., H. Sun, R. M. Polhill, & M. G. Gilbert in Fl. Chin. 10: 107. 2010; Huang T.-C., Ohashi, H. in Fl. Taiwan (2nd Edition). 3: 228-232. 1993. Syn. Crotalaria anagyroides Kunth, Nov. Gen. Sp. (folio ed.) 6: 317. 1824 (Leguminosae).

Caracas rattlebox (三尖叶猪屎豆), originated from C and S America, has been introduced into many tropical countries, including those in Asia and Africa where it has been locally naturalized. It is also cultivated and naturalized in C and S China: Fujian, Guangdong, Guangxi, Hainan, Hubei, Taiwan and Yunnan but sometimes treated as *Crotalaria anagyroides* (Fan *et al.*, 2008).

4. *Duranta erecta* Linnaeus, Sp. Pl. **2**: 637. 1753. Chen S. L. & M. G. Gilbert in Fl. Chin. **17**: 2. 1994. Syn. *Duranta repens* Linnaeus, Sp. Pl. **2**: 637. 1753 (Verbenaceae).

Golden dewdrops (假连翘), native to Tropical America, usually cultivated for ornamental purposes, occasionally escaped and naturalized near farms in Fujian, Guangdong, Guangxi, Hainan,

Hunan, Jiangxi, Macao, Taiwan, Hongkong, Yunnan, Zhengjiang. In some literatures, it was treated as *Duranta repens* (萧如英, 1980; Sun and Xiang, 2004; Ou and Lu, 2006; 解焱, 2008; 陈运造, 2006).

5. Dysphania schraderiana (Schultes) Mosyakin & Clemants, Ukrajins'k. Bot. Žurn. 59: 383. 2002. Zhu G. L., S. L. Mosyakin & S. E. Clemants in Fl. Chin. 5: 377. 2003. Basionym Chenopodium schraderianum Roemer & Schultes, Syst. Veg. 6: 260. 1820. Syn. Chenopodium foetidum Schrader, Mag. Neuesten Entdeck. Gesammten Naturk. Ges. Naturf. Freunde Berlin 2: 79. 1808 (not Lamarck 1778) (Chenopodiaceae).

Foetid Goosefoot (菊叶香藜), species with a strong odor from Africa, SW Asia, and S. Europe but naturalizes in North America and locally elsewhere. In China, it's found in Beijing, Gansu, Liaoning, Nei Mongol, Qinghai, Shaanxi, Shanxi, Sichuan, Xizang, Yunnan, but treated as *Chenopodium foetidum* (孔宪武和简焯坡, 1979; Lin *et al.*, 2009).

6. Eclipta prostrata (Linnaeus) Linnaeus, Mant. Pl. 2: 286. 1771. Chen Y. S. & D. J. Nicholas Hind in Fl. Chin. 20–21: 1481. 2011; Chung K. F. in Fl. Taiwan (2nd Edition). 4: 935 – 937. 1998. Basionym Verbesina prostrata Linnaeus, Sp. Pl. 2: 902. 1753. Syn. Eclipta zippeliana Blume, Bijdr. Fl. Ned. Ind. 914. 1826 (Asteraceae).

False daisy (鳢肠), native to Americas, introduced and naturalized in Africa, Asia, Australia, Europe and Pacific islands. In China, it has spreaded almost all over the country from riversides, fields to abandoned ponds. *Eclipta prostrata* is a highly variable species widely distributed in separated parts of the world and is greatly influenced by the habitats in which they grow. Almost all literatures treated *Eclipta zippeliana* as synonymy of *Eclipta prostrata*, but other treated it as another species (陈运造, 2006).

7. Erigeron bonariensis Linnaeus, Sp. Pl. 2: 863. 1753. Chen Y. L. & L. Brouillet in Fl. Chin. 20 – 21: 1107. 2011. Syn. Conyza bonariensis (Linnaeus) Cronquist, Bull. Torrey Bot. Club 70 (6): 632. 1943; Conyza crispa (Pourret) Ruprecht, Bull. Cl Phys. -Math. Acad. Imp. Sci. Saint-Pétersbourg 14: 235. 1856; Erigeron crispus Pourret, Hist. & Mém. Acad. Roy. Sci. Toulouse 3: 318–319. 1788 (Asteraceae).

Hairy fleabane (香丝草), native to South America, widely naturalized as a weed in tropical and subtropical regions worldwide, including in New Zealand, South Africa, Keny, Tanzania, the USA, Europe and China. It is a kind of crop weeds, particularly serious in no-till farming systems. It has been treated as Conyza bonariensis in almost all Chinese references (Du et al., 2002; Guo et al., 2002; Chen et al., 2003; Wu and Qiang, 2003; Wang et al., 2004; Wu et al., 2004; Hu et al., 2005; Yi et al., 2005; Chen et al., 2006; Zang et al., 2006; Hu et al., 2007; Wang et al., 2007; Li et al., 2008; Tang et al., 2008; Ji et al., 2009; Shu et al., 2009; Wang et al., 2009; Qi et al., 2010; Wang, 2010; 徐海根和强胜, 2011; Ju et al., 2011; Li et al., 2011; Yu et al., 2011; 万方浩等, 2012; Lin et al., 2012b; Zhu, 2012), with the Chinese name as ye tang hao (野塘蒿). Meanwhile, a few literatures used as Conyza crispus (Zhang, 2004; Zhang, 2006; Xie and Zhang, 2012), a wrong spelling of Conyza crispa.

8. Euphorbia hypericifolia Linnaeus, Sp. Pl. 1: 454. 1753. J. S. Ma & Michael G. Gilbert in Fl. Chin. 11: 293. 2008; The Editorial Committee of the Flora of Taiwan in Fl. Taiwan (1st Edition) 6: 79. 1979. Syn. Chamaesyce hypericifolia (Linnaeus) Millspaugh, Publ. Field Columb. Mus., Bot. Ser. 2 (7): 302. 1909 (Euphorbiaceae).

Graceful sandmat (通奶草), native to New World and naturalized in many parts of the Old World. In China it distributes at Anhui, Beijing, Chongqing, Guangdong, Guangxi, Guizhou, Hainan, Hebei, Hongkong, Hubei, Hunan, Jiangxi, Liaoning, Macao, Nei Mongol, Shanxi, Shanghai, Sichuan, Taiwan, Yunnan and Zhejiang. This species has been much confused with *Euphorbia indica* (Liu *et al.*, 2002; Lei *et al.*, 2006) which was called as nan ya da ji (南亚大戟) in Chinese in FOC, but different through the puberulent on stems and capsules (Raju and Rao, 1979).

Euphorbia nutans Lagasca, Gen. Sp. Pl. 17. 1816.
 Shishkin B. K. & E. G. Bobrov in Fl. U. S. S. R. (English edition translated from Russian), 14: 371.
 1974. Syn. Euphorbia preslii Gussone, Fl. Sic. Prodr. 1: 539. 1827 (Euphorbiaceae).

Eyebane (大地锦), a weed in cultivated fields and along roadsides, often along seashores, originated from tropical South America, was introduced into Central Europe and Atlantic Europe and Western Mediterranean. In China, it was distributed in Anhui, Beijing, Hebei, Jiangsu, Liaoning, Shanghai. In some literatures it is treated as *Euphorbia preslii* (Hu *et al.*, 2005).

10. Galinsoga quadriradiata Ruiz & Pavón, Syst. Veg. Fl. Peruv. Chil. 1: 198. 1798. Chen Y. S. & D. J. Nicholas Hind in Fl. Chin. 20 - 21: 1474.
2011. Peng C.-I in Fl. Taiwan (2nd Edition) 4: 968. 1998. Syn. Galinsoga ciliata (Rafinesque)
S. F. Blake, New Fl. 1: 67. 1836 (Asteraceae).

Shaggy soldier (粗毛牛膝菊), species native to tropical America, now has been a cosmopolitan weed. In China, it distributes in Anhui, Chongqing, Guizhou, Hubei, Heilongjiang, Jiangsu, Jiangsi, Liaoning, Shaanxi, Shanghai, Yunnan, Zhejiang and naturlized in northern and central parts of Taiwan along roadsides. In some literatures, it is treated as *Galinsoga ciliata* (Chen et al., 2003; Li et al., 2006; Gao et al., 2008; Mei et al., 2009; Shu et al., 2009; Zhang and Zhang, 2009; Guo et al., 2010; 丁炳扬和胡仁勇, 2011; Yang et al., 2011).

11. *Justicia adhatoda* Linnaeus, Sp. Pl. **1**: 15. 1753; Hu J. Q., Deng Y. F. & T. F. Daniel in Fl. Chin. **19**: 456. 2011. Syn. *Adhatoda vasica* Nees, Pl. Asiat. Rar. **3**: 103. 1832 (Acanthaceae).

Malabar nut (鸭嘴花), probably native to India, Indonesia, Malaysia, Nepal, Pakistan and Sri Lanka, but widely cultivated and naturalized in tropics, and widely used in Ayurvedic and Unani systems of medicine. It has been cultivated and naturalized in Guangdong, Guangxi, Hong Kong, Hainan, Macao, and Yunnan, espeially in the greenhouse for ornamental purpose. In some literatures, it has been treated as Adhatoda vasica (万方浩等, 2012).

12. Nerium oleander Linnaeus, Sp. Pl. 1: 209. 1753. Li P. T., A. J. M. Leeuwenberg & D. J. Middleton in Fl. Chin. 16: 173. 1995. Syn. Nerium indicum Miller, Gard. Dict. (ed. 8) no. 2. 1768 (Apocynaceae).

Oleander (夹竹桃), native to Mediterranean area but now cultivated and often naturalized throughout the tropics, subtropics and temperate parts worldwide (Smith, 1988). In China, it is widely cultivated: Anhui, Beijing, Chongqing, Fujian, Gansu, Guangdong, Guangxi, Guizhou, Hainan, Hebei, Henan, Hongkong, Hubei, Hunan, Jilin, Jiangsu, Jiangxi, Liaoning, Macao, Shaanxi, Shandong, Shanghai, Sichuan, Tianjin, Yunnan, Zhejiang. In Chinese literatures, it has been widely used as Nerium indicum (Fan et al., 2008; Teng, 2008; Sun et al., 2009; Hu et al., 2011). In FRPS, Nerium oleander is treated as ou zhou jia zhu tao (欧洲夹竹桃), with corona not spilited; however, in FOC, Nerium indicum is treated as synonym of Nerium oleander.

Oenothera glazioviana Micheli, Fl. Bras. 13
 13: 178. 1875. Chen J. R., P. C. Hoch & W. L. Wagner in Fl. Chin. 13: 424. 2007. Syn. Oenothera erythrosepala (Bor-bás) Borbás, Magyar Bot. Lapok
 2: 245. 1903 (Onagraceae).

Red sepal evening primrose (黄花月见草),

apparently hybrid origin from the gardens in Europe between two N American species (Raven and Peng, 1993). It was introduced into the horticulture as early as 1860, grown for its particularly large, attractive flowers, and now widely naturalized in Asia, Africa, Europe, Australia, N and S America, Pacific islands (e.g. New Zealand, Hawaii) and in China: Anhui, Beijing, Chongqing, Fujian, Guangdong, Guangxi, Guizhou, Hebei, Henan, Heilongjiang, Hubei, Jilin, Jiangsu, Jiangxi, Liaoning, Nei Mongol, Shaanxi, Shandong, Shanxi, Shanghai, Sichuan, Taiwan, Tianjin, Yunnan, Zhejiang. It's usually called: yue jian cao (月见草) in Chinese, but was treated as Oenothera erythrosepala (Liu et al., 2002; Chen et al., 2003; Yi et al., 2005; 龙茹等, 2008; He and Ge, 2008; Luo and Liu, 2008; Zhou et al., 2008: Li et al., 2009: Jiang et al., 2009: Oi et al., 2010; Zhao et al., 2010; Shi and Li, 2011; Yu et al., 2011; Lin et al., 2012; Zhu and Yang, 2012).

14. Sphagneticola trilobata (Linnaeus) Pruski, Mem. New York Bot. Gard. 87: 114. 1996. J. L. Strother in Fl. N. Amer. 21: 126. 2006. Chen Y. S. & D. J. Nicholas Hind in Fl. Chin. 20-21: 870-871. Basionym Silphium triobatum Linnaeus, Syst. Nat. (ed. 10) 2: 1233. 1759. Syn. Wedelia trilobata (Linnaeus) Hitchcock, Rep. (Annual) Missouri Bot. Gard. 4: 99. 1893 (Asteraceae).

Bay Biscayne creeping-oxeye (南美蟛蜞菊), probably native to subtropics and tropics of the New World and introduced into the Old World. It has been cultivated as ornamental ground cover but escaped in many countries, occasionally naturalized on slopes and along roadsides at lower elevations. It is cultivated and naturalized in Fujian, Guangdong, Guangxi, Hainan, Hongkong, Macao, Sichuan, Taiwan, Yunnna, Zhejiang and usually treated as Wedelia trilobata (李振宇和解焱, 2002; Huang and Zeng, 2003; Zeng, 2003; 徐海根和强胜, 2004; Deng and Lu, 2004; Yan et al., 2006; Guan et al., 2006; Yan et al., 2006; An et al., 2007; Xie et al.,

2007; 解焱, 2008; Chen et al., 2008; Jiang et al., 2008; Wang et al., 2008; Shi et al., 2009; Wang et al., 2009; Yang and Chen, 2009; Zeng et al., 2009; Lin et al., 2011; Yue et al., 2011; Zhang and Xing, 2011; He et al., 2012; Ma et al., 2012; Wu et al., 2012).

15. Stylosanthes guianensis (Aublet) Swartz, Kongl. Vetensk. Acad. Nya Handl. 10: 301–302. 1789. Sa R. & A. D. Salinas in Fl. Chin. 10: 135. 2010. Basionym Trifolium guianense Aublet, Hist. Pl. Guiane 2: 776–778, pl. 309. 1775. Syn. Stylosanthes gracilis Kunth, Nov. Gen. Sp. (quarto ed.) 6: 507–508, pl. 596. 1823 (Leguminosae).

Stylo (圭亚那笔花豆), native to Central and South America, now widely introduced into other tropical areas and sometimes naturalized. It has been cultivated and naturalized in Guangdong, Guangxi, Hainan, Hong Kong and Taiwan. In the past, this species has been used for soil stabilization, however, due to its invasive nature, it has not been recommended for further use. It's also treated as Stylosanthes gracilis in China (Shan et al., 2006; Fan et al., 2008) as ba xi mu xu (巴西苜蓿) in Chinese.

16. Tradescantia pallida (J. N. Rose) D. R. Hunt, Kew Bull. 30 (3): 452. 1975. R. B. Faden in Fl. N. Amer. 22: 173. 2000. Basionym Setcreasea pallida Rose, Contr. U. S. Natl. Herb. 13 (9): 294. 1911. Syn. Setcreasea purpurea Boom, Acta Bot. Neerl. 4: 167. 1955 (Commelinaceae).

Purple queen (紫竹梅), native to Mexico, is widely used as an ornamental plant in gardens and borders including China, as well as a ground cover or hanging pot, since it is remarkably shade-tolerant and can be thrived in a wide range of soil conditions. In southern United States and Australia, it is considered as an invasive species and has defied many attempts at control or eradication. In some literatures, it is also treated as *Setcreasea purpurea* (Chen *et al.*, 2006; Zhang and Zhang, 2009) in China.

Reference ·

- 陈运造,2006. 苗栗地区重要外来入侵植物图志 [M]. 苗栗: 行政院农业委员会苗栗区农业改良场,17,22
- 丁炳扬, 胡仁勇, 2011. 温州外来入侵植物及其研究 [M]. 杭州: 浙江科学技术出版社, 114, 120
- 孔宪武, 简焯坡, 1979. 中国植物志 [M]. 北京: 科学出版社, 25 (2): 80
- 李振宇, 解焱, 2002. 中国外来入侵种 [M]. 北京: 中国林业出版社, 174
- 龙茹, 史风玉, 孟宪东等, 2008. 河北省外来入侵植物调查分析 [J]. 北方园艺, 7: 171—173
- 曲波, 2003. 沈阳地区外来入侵有害植物的调查 [J]. 辽宁农业科学, **4**: 29—31
- 万方浩, 刘全儒, 谢明等, 2012. 生物人侵: 中国外来人侵植物图鉴 [M]. 北京: 科学出版社.
- 吴桂容, 2006. 广西外来入侵植物的初步研究 [J]. 广西梧州师 范高等专科学校学报, 22 (2): 96—100
- 解焱,2008.生物入侵与中国生态安全[M].石家庄:河北科学技术出版社,336,440
- 徐海根,强胜,2004. 中国外来入侵物种编目 [M]. 北京:中国 环境科学出版社,183
- 徐海根,强胜,2011. 中国外来入侵生物 [M]. 北京: 科学出版 社,283—284
- 萧如英,1980. 台湾植物志 (第一版) [M]. 台北: 现代关系出版社,4:425
- An F (安锋), Kan LY (阚丽艳), Xie GS (谢贵水) et al., 2007.

 Alien invasive plants in Hainan island and contral countermeasures [J]. Journal of Northwest Forestry University (西北林学院学报), 22 (5): 198—206
- Chen J (陈京), Xu P (徐攀), Yao ZS (姚振生), 2011. Analysis and control of invasive plants of Tianmushan nature reserve [J]. China Medical Herald (中国医学导报), **8** (9): 137—139
- Chen ML (陈明林), Zhang XP (张小平), Su DS (苏登山), 2003. A preliminary study on the invasive weeds in Anhui [J]. *Journal of Biology* (生物学杂志), **20** (6): 24—27
- Chen QX (陈秋霞), Wei CQ (韦春强), Tang SC (唐赛春) et al., 2008. The alien invasive plants in Guilin city, Guangxi Zhuangzu autonomous region [J]. Subtropical Plant Science (亚热带植物科学), 37 (3): 55—58
- Chen XY (陈小永), Wang HY (王海燕), Ding BY (丁炳扬) et al., 2006. The species and habitat characteristics of exotic weeds in Hangzhou [J]. Bulletin of Botanical Research (植物研究), 26 (2): 242—249
- Deng XC (邓晰朝), Lu X (卢旭), 2004. A preliminary study on the invasive plants in Yizhou city [J]. *Journal of Hechi Normal College* (河池师专学报), **24** (2): 72—74
- Du WB (杜卫兵), Ye YZ (叶永忠), Zhang XY (张秀艳) et al., 2002. Study on the foreing harmful plant in Henan [J]. Henan Science (河南科学), 20 (1): 52—55
- Fan ZW (范志伟), Shen YD (沈奕德), Liu LZ (刘丽珍), 2008.

- List of invasive alien weeds in Hainan, China [J]. Chinese Journal of Tropical Crops (热带作物学报), **29** (6): 781—792
- Fu HB (付海滨), Zhang M (张敏), Kang K (康凯) et al., 2009.

 Main alien invasive species of agriculture and forestry and control strategy in Shenyang [J]. Environmental Protection Science (环境保护科学), 35 (3): 31—33
- Fu L (付岚), Zhao MF (赵鸣飞), Gong L (龚玲) et al., 2012.

 Investigation and risk assessment of alien invasive plants in riparian zone of Dongjiang river [J]. Journal of Anhui Agricultural Sciences (安徽农业科学), 40 (3): 1689—1693, 1788
- Gao M (高末), Luo ZR (骆争荣), Ge LQ (戈丽清) et al., 2008.

 Distribution and spread characteristics of invasive plants of compositae in Wenzhou [J]. Journal of Wenzhou University (Natural Sciences) (温州大学学报:自然科学版), 29 (5):17—24
- Gao Y (高燕), Cao W (曹伟), 2010. Situation and countermeasure of alien invasive plants in Northeast China [J]. Journal of the Graduate School of the Chinese Academy of Sciences (中国科学院研究生院学报), 27 (2): 191—198
- Guan ZB (管志斌), Deng WH (邓文华), Huang ZL (黄志玲) et al., 2006. A preliminary investigation on the alien invasive plants in Xishuangbanna [J]. Tropical Agricultural Science & Technology (热带农业科技), 29 (4): 35—38
- Guo SL (郭水良), Mao YR (毛郁薷), Qiang S (强胜), 2002. Influences of temperature on peroxidase isozymoframs of six exotic weeds [J]. Guihaia (广西植物), 22 (6): 557—562
- Guo YQ (郭怡卿), Zhao GJ (赵国晶), Chen Y (陈勇) et al., 2010. Infestation of invasive weed in crop field in Yunnan [J]. Southwest China Journal of Agricultural Sciences (西南农业学报), 23 (4): 1352—1355
- He JQ (何家庆), Ge JL (葛结林), 2008. The present situation of alien invasive plants in Anhui province and a comparison with the other provinces in China [J]. *Journal of Anhui University* (Natural Science Edition) (安徽大学学报:自然科学版), **32** (2): 82—89
- He TP (何太平), Hu G (胡刚), Zhang ZH (张忠华), 2012.
 Alien invasive plants in Nanning, China [J]. Journal of Tropical and Subtropical Botany (热带亚热带植物学报), 20 (5): 497—505
- Hu G (胡刚), Zhang ZH (张忠华), Liang SC (梁士楚), 2005.

 Preliminary study on the alien weeds in Huaibei area [J].

 Journal of Anhui Agricultural Sciences (安徽农业科学), 33
 (5): 789—790, 819
- Hu TY (胡天印), Huang H (黄华), Guo SL (郭水良) *et al.*, 2007. Niche characteristic of exotic weeds in suburb and the impacts on biodiversity [J]. *Guihaia* (广西植物), **27** (6): 873—881
- Hu XH (胡雪华), Xiao YA (肖宜安), Zeng JJ (曾建军) et al., 2011. Investigation and analysis on alien plants in Bawangling national nature reserve in Hainan Province [J]. Journal of Jinggangshang University (Natural Science) (井冈山大学学

- 报: 自然科学版), 32(1): 131-136
- Huang CP (黄彩萍), Zeng LM (曾丽梅), 2003. The investigation on the harm and prevention strategies of the exot invasion weeds in the Guangzhou white cloud mountain [J]. *Tropical Forestry* (热带林业), **31** (4): 24—29
- Ji CF (季春峰), Wang Z (王智), Qian P (钱萍), 2009. Primary research on invasive plants in Jiangxi province [J]. *Hubei Forest-ry Science and Technology* (湖北林业科技), **3**: 27—31
- Jiang GC (姜罡丞), Wang K (王珂), Dong DP (董东平), 2009.

 Research and risk assessment of alien invasive plants of Xuchang suburb [J]. Journal of Henan Normal University (Natural Science) (河南师范大学学报: 自然科学版), 37 (6): 168—170
- Jiang QC (蒋谦才), Lin ZM (林正眉), Li L (李荔) et al., 2008.

 An investigation on the invasive plants in Zhongshan, Guangdong Province [J]. Guangdong Forestry Science and Technology (广东林业科技), 24 (2): 54—58
- Ju JW (鞠建文), Wang N (王宁), Guo YJ (郭永久) et al., 2011. Current situation analysis of invasive plant in Jiangxi province [J]. Journal of Jinggangshan University (Natural Science) (井冈山大学学报:自然科学版), 32 (1): 126—130
- Li B (李斌), Wang Y (王咏), He CG (何春光), 2007. Primary Research on exotic plants in Changchun area [J]. *Journal of Changchun Normal University* (Natural Science) (长春师范学院学报: 自然科学版), **26** (5): 85—88
- Li GY (李根有), Jin SH (金水虎), Ai JG (哀建国), 2006. Species, characteristics and control measures of injurious plants in Zhejiang province [J]. Journal of Zhejiang Forestry College (浙江林学院学报), 23 (6); 614—624
- Li SC (李顺才), Liu LX (刘连新), Hu XD (胡晓东) et al., 2009. Investigation and analysis of alien invasions in Qinhuangdao [J]. Journal of Hebei Normal University of Science & Technology (河北科技师范学院学报), 23 (3): 13—17
- Li Y (李亚), Yao G (姚淦), Deng F (邓飞) et al., 2008. Preliminary investigation and analysis on exotic spermatophyte in Jiangsu province [J]. Journal of Plant Resources and Environment (植物资源与环境学报), 17 (4): 55—60
- Li YM (李玉梅), Lu SF (陆山风) *et al.*, 2011. Investigation and study on the alien invasive plants along nan-you expressway of Guangxi [J]. *Guangxi Forestry Science* (广西林业科学), **40** (4): 277—280
- Lei T (雷霆), Ding LJ (丁立建), Zhang JR (张佳蕊) et al., 2006. Flora characteristics of vascular plants in wetland nature reserves in Beijing [J]. Journal of Beijing Forestry University (北京林业大学学报), 28 (6): 67—74
- Lin JY (林建勇), Wen YG (温远光), Wei J (韦洁), 2011. The alien invasive plants in Guangxi beibu gulf economic zone [J].

 Guangxi Forestry Science (广西林业科学), 40 (4): 281—287
- Lin JY (林建勇), Liang RL (梁瑞龙), Li J (李娟) et al., 2012a.

 Study on the invasive alien plants in south China [J]. Guangxi Forestry Science (广西林业科学), 41(3); 237—241

- Lin M (林敏), Hao JH (郝建华), Chen GQ (陈国奇), 2012b.

 Composition and distribution of invasive alien plants in Suzhou area [J]. Journal of Plant Resources and Environment (植物资源与环境学报), 21 (3): 98—104
- Lin QW (林秦文), Xing SH (邢韶华), Ma K (马坤), 2009. New materials of alien plants in Beijing [J]. *Journal of Beijing University of Agriculture* (北京农学院学报), **24** (4): 42—44
- Liu QR (刘全儒), Yu M (于明), Zhou YL (周云龙), 2002. A preliminary study on the invasive plants in Beijing [J]. *Journal of Beijing Normal University* (Natural Science) (北京师范大学学报: 自然科学版), **38** (3): 399—404
- Luo Y (罗艳), Liu AH (刘爱华), 2008. Preliminary research on alien invasive plants of Qingdao [J]. Shandong Science (山东科学), 21 (4): 19—23
- Ma D (马多), He TP (和太平), Zheng X (郑羨), 2012. Study on alien invasive plants in Wuzhou city [J]. Guangxi Forestry Science (广西林业科学), 41 (2): 155—158
- Ma JS (马金双), 2013. The Checklist of the Chinese Ivasive Plants [M]. Beijing: High Education Press
- Mei XM (梅笑漫), Ding BY (丁炳扬), Jin XF (金孝锋), 2009. Study on alien weeds of the west-iake scenic spot in Hangzhou [J]. Guihaia (广西植物), **29** (1): 125—131
- Qi YZ (祁云枝), Du YJ (杜勇军), Zhang Y (张莹), 2010. A investigation of alien invasive plants in Xian area [J]. Chinese Agricultural Science Bulletin (中国农学通报), 26 (5): 223—227
- Ou J (欧健), Lu CY (卢昌义). 2006. The research of alien plants risk assessment system in Xiamen municipality [J]. Journal of Xiamen University (Natural Science) (厦门大学学报:自然科学版), 45 (6): 883—888
- Qu B (曲波), Lv GZ (吕国忠), Yang H (杨红) et al., 2006a. Report about alien invasive plants in Liaoning province [J]. Journal of Shenyng Agricultural University (沈阳农业大学学报), 37 (4): 587—592
- Qu B (曲波), Lv GZ (呂国忠), Yang H (杨红) et al., 2006b.

 Preliminary report about alien invasive plants in Liaoning province [J]. Liaoning Agricultural Sciences (辽宁农业科学), 4: 22—25
- Raven, Peter H, Chin-I Peng, 1993. Fl. Taiwan (2nd Edition)
 [M]. Taipei; Lungwei Printing Company, Ltd., 3: 965
- Shan JL (单家林), Yang FC (杨逢春), Zheng XQ (郑学勤), 2006. Exotic plants in Hainan province [J]. Subtropical Plant Science (亚热带植物科学), 35 (3): 39—44
- Raju VS, Rao PN, 1979. E. indica: hitherto little known name of a much common Indian weed [J]. Indian Journal of Botany, 2 (2): 202—208
- Shi DH (石登红), Li C (李灿), 2011. Alien plant invasion situation in "two lakes and one reservoir" ecological functional area of Guiyang and prevention measures [J]. Guizhou Agricultural Sciences (贵州农业科学), 39 (3): 94—97
- Shi LC (石亮成), Shi G (石钢), Yi QL (易巧玲) et al., 2009.

- The countermeasure research for investigation and eradication of alien invasive plant in Liuzhou [J]. *Journal of Guangxi Academy of Sciences* (广西科学院学报), **25** (3): 178—182
- Shu MY (舒美英), Cai JG (蔡建国), Fang BS (方宝生), 2009.

 Situation and control measure of alien invasive plants in Xixi national wetland park, Hangzhou city, Zhejiang province [J].

 Journal of Zhejiang Forestry College (浙江林学院学报), 26
 (5): 755—761
- Smith, Albert C, 1988. Flora Vitiensis Nova: a New Flora of Fiji M]. Lawaii, Hawaii; Pacific Tropics Botanical Garden, 4: 193
- Sun J (孙娟), Yang GF (杨国锋), Chen YC (陈玉成) et al., 2009. Exotic plants in the Jinfo Mountain Nature Reserve: species ascertain and their distribution [J]. Acta Prataculturae Sinica (草业学报), 18 (3): 34—42
- Sun WB (孙卫邦), Xiang QB (向其柏), 2004. Introduction and utilization of the alien ornamentals and biotic invasion [J]. *Chinese Landscape Architecture* (中国园林), **20** (9): 54—56
- Tang SC (唐赛春), Lv SH (吕仕洪), He CX (何成新) et al., 2008. The alien invasive plants in Guangxi [J]. Guihaia (广西植物), 28 (6): 775—779
- Teng YQ (滕永清), 2008. Exotic plant species in Jinfoshan nature reserve [D]. Chongqing; Southwest University
- Wang F (王芳), Wang RJ (王瑞江), Zhuang PD (庄平弟) et al., 2009. Present status and anagement strategies of alien invasive plants in Guangdong province [J]. Chinese Journal of Ecology (生态学杂志), 28 (10): 2088—2093
- Wang FG (王发国), Xing FW (邢福武), Ye HG (叶华谷) et al., 2004. Preliminarily study on invasive species in Macau [J]. Acta Scientiarum Naturalium Universitatis (Sunyatseni) (中山大学学报:自然科学版), 43 (Suppl.): 106—110
- Wang SM (王苏铭), Zhang N (张楠), Yu LQ (于琳倩) et al., 2012. Distribution pattern and their influcing factors of invasive alien plants in Beijing [J]. Acta Ecologica Sinica (生态学 报), **32** (15): 4618—4629
- Wang W (王伟), Zhang XM (张先敏), Sha LH (沙林华) et al., 2007. Roster of alien invasive perilous species of animals and plants in Hainan island (I) [J]. Chinese Journal of Tropical Agriculture (热带农业科学), 27 (4): 58—64
- Wang YJ (王元军), 2010. Investigation on alien invasive plants in the nansihu lake wetland of Shandong province [J]. *Chinese Bulletin of Botany* (植物学报), **45** (2): 212—219
- Wang Z (王忠), Dong SY (董仕勇), Luo YY (罗燕燕) et al., 2008. Invasive plants in Guangzhou, China [J]. Journal of Tropical and Subtropical Botany (热带亚热带植物学报), 16 (1): 29—38
- Wu HR (吴海荣), Qiang S (强胜), Lin JC (林金成), 2004.

 Quantitative survey on spring exotic weeds and the niche characteristics of dominant species in Nanjing [J]. Acta Botanica Boreali-Occidentalia Sinica (西北植物学报), 24 (11): 2061—2068

- Wu HR (吴海荣), Qiang S (强胜), 2003. Quantitative survey on exotic weeds in autumn in Nanjing [J]. *Biodiversity Science* (生物多样性), **11** (5): 432—438
- Wu J (吴杰), Qin M (覃铭), Zeng XH (曾晓华) et al., 2012. A survey of higher plants in Beijing normal University, Zhuhai [J]. Journal of Beijing Normal University (Natural Science) (北京师范大学学报:自然科学版), 48 (2): 188—194
- Wu YQ (吴彦琼), Hu YJ (胡玉佳), Chen JN (陈江宁), 2005.
 Repreoductive characteristics of alien plant Wedelia trilobata
 [J]. Acta Scientiarum Naturalium Universitatis Sunyatseni (中山大学学报:自然科学版), 44 (6):93—96
- Xie YZ (谢云珍), Wang YB (王玉兵), Tan WF (谭伟福), 2007. Invasive plants in Guangxi province [J]. *Journal of Tropical and Subtropical Botany* (热带亚热带植物学报), **15** (2): 160—167
- Xie HY (谢红艳), Zhang XQ (张雪芹), 2012. Research on invasive alien plant in Hunan province [J]. Modern Agricultural Science and Technology (现代农业科技), 5: 178—179, 181
- Yan SJ (闫淑君), Hong W (洪伟), Wu CZ (吴承祯), 2006. Effects of biological invasion on ecological safety of Fujian province [J]. *Journal of Fujian College of Forestry* (福建林学院学报), **26** (3): 275—280
- Yan YH (严岳鸿), Xing FW (邢福武), Huang XX (黄向旭) et al., 2004. Exotic plants in Shenzhen, China [J]. Guihaia (广西植物), 24 (3): 232—238
- Yang D (杨德), Liu GH (刘光华), Xiao CM (肖长明) et al., 2011. Identification and control of invasive agricultural plants in Chongqing [J]. Acta Agriculturae Jiangxi (江西农业学报), 23 (3): 93—95
- Yang J (杨坚), Chen HB (陈恒彬), 2009. A preliminary study on alien invasive plants in Fujian province [J]. Subtropical Plant Science (亚热带植物科学), 38 (3): 47—52
- Yi YJ (衣艳君), Li XS (李修善), Qiang S (强胜), 2005. Study on exotic weeds in Shandong province [J]. Territory & Natural Resources Study (国土与自然资源研究), 3:87—88
- Yue MF (岳茂峰), Fan BL (樊蓓莉), Tian XS (田兴山) et al., 2011. Investigation and hazard evaluation of exotic invasive plants in agricultural ecosystems in Guangdong probince [J]. Journal of Biosafety (生物安全学报), 20 (2): 141—146
- Yu DZ (喻大昭), Li RH (李儒海), Chu SH (褚世海) et al., 2011. Invasive species in Hubei province, China and its relationship with the social and economic activities [J]. *Journal of*

- Biosafety (生物安全学报), 20 (1): 56—63
- Zang M (臧敏), Qiu XL (邱筱兰), Huang LF (黄立发) et al., 2006. Study on the alien plants in Anhui province [J]. Journal of Anhui Agricultural Sciences (安徽农业科学), **34** (20): 5306—5308
- Zeng XF (曾宪锋), Lin XD (林晓单), Qiu HY (邱贺媛) et al., 2009. Investigation on alien plants in E. Guangdong [J]. Journal of Fujian Forestry Science and Technology (福建林业科技), **36** (2): 174—179, 249
- Zeng XF (曾宪锋), 2003. A preliminary study on 5 species invasive plants in E. Guangdong [J]. *Journal of Hanshan Teachers College* (韩山师范学院学报), **24** (3): 64—69, 71
- Zhang GB (张桂宾), 2004. Study on the Invasive alien plant in Kaifeng [J]. *Journal of Henan University* (Natural Science) (河南大学学报:自然科学版), **34** (1):56—59
- Zhang GB (张桂宾), 2006. Invasive alien plants and control in the farming region of Yudong [J]. *Journal of Shangqiu Teachers College* (商丘师范学院学报), 22 (2): 153—156
- Zhang JG (张建国), Zhang MR (张明如), 2009. Rearch on the establishment of the early-warning system for invasion of injurious biology outside Zhejiang [J]. *Journal of Fujian Forestry Science and Technology* (福建林业科技), **36** (1): 100—105
- Zhang RJ (张荣京), Xing FW (邢福武), 2011. The composition and origin of alien plants in Ganshiling nature reserve in Hainan, China [J]. Guizhou Agricultural Sciences (贵州农业科学), 39 (7): 31—33
- Zhao JJ (赵娟娟), OuYang ZY (欧阳志云), Zheng H (郑华) et al., 2010. Species composition of alien plants in the built-up area of Beijing [J]. Biodiversity Science (生物多样性), 18 (1): 19—28
- Zhou XG (周小刚), Chen QH (陈庆华), Zhang H (张辉) et al., 2008. Invasive alien weeds species in farmland and forest in Sichuan province [J]. Southwest China Journal of Agricultural Sciences (西南农业学报), 21 (3): 852—858
- Zhu BH (朱碧华), Yang FM (杨凤梅), 2012. Alien invasive plants in Nanchang urban green land and its preventive measures [J]. *Jiangxi Forestry Science and Technology* (江西林业科技), 1: 35—38
- Zhu H (朱慧), 2012. Study on the clonality and invasive of invasive plants in east Guangdong [J]. Chinese Agricultural Science Bulletin (中国农学通报), 28 (15): 199—206